



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

FOREIGN STATISTICAL REPORTS.

FRANCE—*Rouen*.—Month of June, 1899. Estimated population, 112,657. Total number of deaths, 240, including enteric fever, 2; scarlet fever, 2; whooping cough, 3, and 50 from phthisis pulmonalis.

GERMANY—*Stuttgart*.—Month of April, 1899. Estimated population, 174,434. Total number of deaths, 163, including diphtheria, 4; measles, 2, and 23 from phthisis pulmonalis.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 33 great towns in England and Wales during the week ended July 15, 1899, correspond to an annual rate of 16.9 a thousand of the aggregate population, which is estimated at 11,404,408. The highest rate was recorded in Manchester, viz, 24.8, and the lowest in Croydon, viz, 10.6.

London.—One thousand three hundred and eighty-seven deaths were registered during the week, including measles, 48; scarlet fever, 4; diphtheria, 37; whooping cough, 26; enteric fever, 6; smallpox, 1, and diarrhea and dysentery, 75. The deaths from all causes correspond to an annual rate of 15.9 a thousand. In Greater London 1,781 deaths were registered, corresponding to an annual rate of 14.2 a thousand of the population. In the "outer ring" the deaths included 13 from diphtheria, 7 from measles, 1 from scarlet fever, and 3 from whooping cough.

Ireland.—The average annual death rate represented by the deaths registered during the week ended July 15, 1899, in the 22 principal town districts of Ireland was 21.2 a thousand of the population, which is estimated at 1,039,480. The lowest rate was recorded in Tralee, viz, 0.0, and the highest in Newtownards, viz, 34.0 a thousand. In Dublin and suburbs 173 deaths were registered, including diphtheria, 3; enteric fever, 1; measles, 7, and whooping cough, 4.

Scotland.—The deaths registered in 8 principal towns during the week ended July 15, 1899, correspond to an annual rate of 16.7 a thousand of the population, which is estimated at 1,587,414. The lowest mortality was recorded in Leith, viz, 11.4, and the highest in Paisley, viz, 21.6 a thousand. The aggregate number of deaths registered from all causes was 510, including diphtheria, 2; measles, 8; scarlet fever, 3, and whooping cough, 12.

ITALY—*Florence*.—Month of April, 1899. Estimated population, 199,080. Total number of deaths, 335, including enteric fever, 2; whooping cough, 4, and 36 from phthisis pulmonalis.

JAMAICA.—Two weeks ended July 15. Estimated population, 639,491. Number of deaths not reported. No contagious diseases reported.